



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Again, $CH + CD = 2CF = 2CE \cdot \frac{CF}{CE} = 2CE \cdot \frac{CK}{CM} = 2CE \cdot CK$,

$$\text{or } \cos x + \cos y = 2 \cos \frac{1}{2}(x+y) \cos \frac{1}{2}(x-y) \dots \dots \dots (2).$$

The triangles CEM and XNY are similar ;

$$\text{hence } \frac{NX}{NY} = \frac{CE}{CM}, \text{ or } NX = 2CE \cdot \frac{XY}{CM} = 2CE \cdot KY,$$

$$\text{that is, } \sin x - \sin y = \cos \frac{1}{2}(x+y) \sin \frac{1}{2}(x-y) \dots \dots \dots (3).$$

Similarly, $\frac{NY}{XY} = \frac{EM}{CM}$, or $NY = 2EM \cdot \frac{XY}{CM} = 2EM \cdot KY$,

$$\text{or } \cos x - \cos y = -2 \sin \frac{1}{2}(x+y) \sin \frac{1}{2}(x-y) \dots \dots \dots (4).$$

Equation (1) can be made very useful in computing trigonometric tables, as the writer intends subsequently to show.

Now let $AM = x$ and $MY = MX = y$. Then $AY = x - y$ and $AX = x + y$.

We have $(CM)^2 - (CK)^2 = (CY)^2 - (CK)^2 = (KY)^2$. But $\frac{CM}{ME} = \frac{CK}{KF} = \frac{KY}{LY}$.

$$\text{Therefore } (ME)^2 - (KF)^2 = (LY)^2 = (KY)^2 - (KL)^2,$$

$$\text{or } (KF)^2 - (KL)^2 = (ME)^2 - (KY)^2,$$

$$\text{or } (KF + KL)(KF - KL) = (ME)^2 - (KY)^2,$$

$$\text{or } HX \times DY = (ME)^2 - (KY)^2.$$

$$\text{That is, } \sin(x+y) \sin(x-y) = \sin^2 x - \sin^2 y = \cos^2 y - \cos^2 x \dots \dots \dots (5).$$

$$\text{Again, } \frac{CM}{CE} = \frac{CK}{CF} = \frac{KY}{KL}.$$

$$\text{Therefore } (CE)^2 - (CF)^2 = (KL)^2 = (KY)^2 - (LY)^2,$$

$$\text{or } (CF)^2 - (LY)^2 = (CE)^2 - (KY)^2,$$

$$\text{or } (CF - LY)(CF + LY) = CH \times CD = (CE)^2 - (KY)^2.$$


$$\text{That is, } \cos(x+y) \cos(x-y) = \cos^2 x - \sin^2 y = \cos^2 y - \sin^2 x \dots \dots \dots (6).$$

Let $DC = R$ the radius of a circle. Let the angle $CDB = 2x$. Then $DAB = DBA = CBE = x$.

$$\text{Then we have } \tan x = \frac{EC}{EB}, \text{ also } \tan x = \frac{BE}{AE}.$$

$$\text{The product of these gives, } \tan^2 x = \frac{CE}{AE}, \text{ or } CE \times AE = (BE)^2,$$

$$\text{or } \frac{EC}{AE} = \left(\frac{BE}{AE} \right)^2 = \tan^2 x.$$



$$\text{Also, } \frac{EC}{BE} = \frac{\text{vers}2x}{\sin2x} = \frac{1-\cos2x}{\sin2x} = \frac{\sin2x}{1+\cos2x} = \tan x \text{ [see above] } \dots\dots\dots(7).$$

$$\text{Then } 1+\tan^2x = 1 + \frac{EC}{AE} = \frac{AC}{AE} = \frac{2R}{AE}$$

$$1-\tan^2x = 1 - \frac{EC}{AE} = \frac{AE-EC}{AE} = \frac{AC-2EC}{AE} = \frac{2(R-EC)}{AE}.$$

$$\cot2x = \frac{DE}{BE} \text{ and } \operatorname{cosec}2x = \frac{R}{BE}. \text{ From these values we at once have,}$$

$$\frac{2\tan x}{1+\tan^2x} = \frac{2BE}{AE} \cdot \frac{AE}{2R} = \frac{BE}{R} = \sin2x \dots\dots\dots(8).$$

$$\frac{2\tan x}{1-\tan^2x} = \frac{2BE}{AE} \cdot \frac{AE}{2(R-EC)} = \frac{BE}{R-EC} = \frac{BE}{DE} = \tan2x \dots\dots\dots(9).$$

$$\tan^2x + 2\cot2x\tan x = \frac{EC}{AE} + \frac{2DE}{BE} \cdot \frac{BE}{AE} = \frac{EC+2DE}{AE} = \frac{AE}{AE} = 1 \dots\dots\dots(10).$$

$$2\operatorname{cosec}2x\tan x - \tan^2x = \frac{2R}{BE} \cdot \frac{BE}{AE} - \frac{EC}{AE} = \frac{2R-EC}{AE} = \frac{AE}{AE} = 1 \dots\dots\dots(11).$$

$$\frac{1-\tan^2x}{1+\tan^2x} = \frac{2(R-EC)}{AE} \cdot \frac{AE}{2R} = \frac{R-EC}{R} = \frac{ED}{R} = \cos2x \dots\dots\dots(12).$$

$$\operatorname{cosec}2x - \cot2x = \frac{R-ED}{BE} = \frac{EC}{BE} = \tan x = \frac{1-\cos2x}{\sin2x} = \frac{\sin2x}{1+\cos2x} \dots\dots\dots(13).$$

$$\operatorname{cosec}2x + \cot2x = \frac{R+ED}{BE} = \frac{AE}{BE} = \cot x = \frac{\sin2x}{1-\cos2x} = \frac{1+\cos2x}{\sin2x} \dots\dots\dots(14).$$

$$\frac{1+\sin2x-\cos2x}{1+\sin2x+\cos2x} = \frac{R+BE-ED}{R} \div \frac{R+BE+ED}{R} = \frac{EC+BE}{AE+BE}.$$

$$\begin{aligned} \text{But } AE &= \frac{(BE)^2}{EC}; \therefore \frac{EC+BE}{AE+BE} = \frac{EC+BE}{(BE)^2 \div EC + BE} \\ &= \frac{EC(EC+BE)}{BE(EC+BE)} = \frac{EC}{BE} = \tan x \dots\dots\dots(15). \end{aligned}$$

Again, $\cos x = \frac{AE}{AB}$, also $\cos x = \frac{AB}{AC}$.

Twice the product of these gives $2\cos^2 x = \frac{2AE}{AC} = \frac{AE}{R}$.


Also $\cos 2x = \frac{DE}{R}$. $1 + \cos 2x = \frac{DE + R}{R} = \frac{AE}{R}$. $\therefore 1 + \cos 2x = 2\cos^2 x \dots \dots (16)$.

$\sin x = \frac{CB}{AC} = \frac{BC}{2R}$; also $\sin x = \frac{EC}{BC}$. Twice the product of these gives

$2\sin^2 x = \frac{EC}{R}$. $1 - \cos 2x = \frac{R - ED}{R} = \frac{EC}{R}$. $\therefore 1 - \cos 2x = 2\sin^2 x \dots \dots \dots (17)$.

To prove the "Tangent Proportion," let ABC be a plane triangle, the parts being represented as usual. Take $CE = CA$ and draw AEH . Draw BHK perpendicular to AH , to meet AC prolonged in K . Now considering the triangles ABC and ACE , the sum of the angles at A and E of the one is equal to the sum of the angles at A and B of the other. Hence $CAE + CEA = A + B$; and $CAE = CEA = BEH = \frac{1}{2}(A + B)$.

Also $BAE = A - \frac{1}{2}(A + B) = \frac{1}{2}(A - B)$. The angles at B and K of the triangle BCK are equal; for CBK is the complement of BEH or AEC , and BKC is the complement of the equal angle CAE . Hence $CK = CB = a$ and $AK = a + b$.



Now $\tan \frac{1}{2}(A - B) = \frac{BH}{AH}$ and $\tan \frac{1}{2}(A + B) = \frac{HK}{AH}$. $\therefore \frac{\tan \frac{1}{2}(A - B)}{\tan \frac{1}{2}(A + B)} = \frac{BH}{HK}$.

But $\frac{BH}{HK} = \frac{BE}{AK} = \frac{a - b}{a + b}$. $\therefore \frac{\tan \frac{1}{2}(A - B)}{\tan \frac{1}{2}(A + B)} = \frac{a - b}{a + b} \dots \dots \dots (1)$.

From the triangle ABE , $\frac{BE}{AB} = \frac{\sin BAE}{\sin AEC}$, or $\frac{a - b}{c} = \frac{\sin \frac{1}{2}(A - B)}{\sin \frac{1}{2}(A + B)} \dots (2)$.

In the triangle AHK , $AH = AK \cos HAK = (a + b) \cos \frac{1}{2}(A + B)$.

In the triangle ABH , $AH = AB \cos BAH = c \cos \frac{1}{2}(A - B)$.

Equating, we have, $\frac{a + b}{c} = \frac{\cos \frac{1}{2}(A - B)}{\cos \frac{1}{2}(A + B)} \dots \dots \dots (3)$.

Equation (3) divided by (2) also gives (1).